

## **Multiple Psychiatric Medications in the Treatment of Child and Adolescent Psychiatric Disorders**

Recent surveys of psychotropic drug usage over the last decade report an increase in polypharmacy in youth with psychiatric disorders. There are many factors that can lead to the use of multiple medications in children and adolescents. While treatment decisions should always reflect the results of evidence-based best practice guidelines, clinicians treating youth with psychiatric disorders find disappointingly little guidance from the psychiatric literature. There are only a small number of either well-designed and useful studies completed on youth or comprehensive practice guidelines from the American Academy of Pediatrics and the American Academy of Child and Adolescent Psychiatry. Additionally, many of the older psychotropic drugs in common usage with children and adolescents have not had the stringent clinical trials that the FDA currently requires.

### **Practice Challenges**

Significant practice challenges are posed by limited scientific evidence and few consensus guidelines for treating children and adolescents with psychiatric disorders. This situation is further complicated by the need to distinguish symptoms of psychiatric disorders from developmental changes, and to respond to the pleas of concerned teachers and parents. It is crucial to obtain input from parents as part of the evaluation. It may also be important to explain that there may not be a “quick fix” for psychiatric symptoms, and that developmental changes complicate determination of diagnosis. Often, practitioners undertake evaluation and treatment in a context of crisis and decreased hospital lengths of stay. Medication trials are one of a few reasons why insurance companies will approve additional hospital days. However, many practitioners attempt to treat psychiatric symptoms in youth by prescribing medications for which there is no clear evidence-based support or expert consensus recommendation. In these cases, achieving the ideal, meticulous, single-drug medication trial is even more difficult. Inappropriate medication usage, as well as the use of multiple medications may result.

Despite these challenges, the use of multiple medications may be appropriate in the treatment of some children and adolescents with complex psychiatric disorders. For example, if a child has several psychiatric diagnoses that are uniquely responsive to different classes of medications, a clinician may prescribe several medications (e.g., a child with Tourette’s syndrome, OCD and ADHD may require a psychostimulant, a SSRI antidepressant and, possibly another medication).

Jellinek cites, with caution, the benefit of combined pharmacotherapy in its effective reduction of psychiatric symptoms that allows for other interventions to take hold. For example, a child with severe OCD whose symptoms are only partially responsive to fluvoxamine and cognitive-behavioral therapy may benefit more fully from the addition of a small dose of clomipramine.

The addition of another drug, to treat the side effects of intervention, is appropriate and the preferred method of treatment. For example, a child with psychosis who is treated with risperidone and develops severe EPS may be prescribed an anti-cholinergic to

relieve the EPS symptoms. An example of treatment with an older drug not subjected to current prevailing FDA standards is the use of trazodone as a hypnotic.

The National Association of State Mental Health Program Directors' (NASMHPD) has identified the following risks inherently associated with polypharmacy:

- ❑ The use of multiple medications increases the risk for medication-related adverse events and drug interactions.
- ❑ The use of multiple medications creates a more complicated drug regimen for the patient, potentially making adherence more difficult.
- ❑ Multiple medications may confound the effects of one another.
- ❑ Where medications are used to treat the side effects of other medications, polypharmacy potentially creates the need for more medication.
- ❑ The medication costs must be borne by the patient or another payer.

### **Recommendations:**

Given the challenges to the effective prescription of psychotropic medication that have been described above, the following recommendations may help practitioners to avoid unnecessary and/or dangerous polypharmacy:

1. Do not rely on anecdotal reports, “word of mouth” experiences, but instead always consider available evidence. Double-blind placebo controlled clinical studies are the gold standard and should be considered primarily. When empirical evidence is lacking, an alternative source for guidance comes from the consensus of experts.
2. Psychotropic medications should only be prescribed after a thorough biopsychosocial evaluation. In complex cases, evaluation should include medical and neurological exams, as well as psychological and psychiatric assessments. Psychotropic medications should be targeted at specific symptoms that are related to specific DSM-IV diagnoses.
3. Collaborative treatment planning and education with the patient, parents and/or guardians is essential to ensuring effective treatment, promoting adherence to the treatment regimen and effecting reliability in reporting on persistent symptoms, side effects and significant events to guide continuing treatment.
4. “Medication only” treatment is usually not appropriate, especially in the early stages of intervention. Addition of effective non-medication interventions is indicated for maximum efficacy and improvement. Examples of effective non-medication interventions are Parent Management Training for oppositional behavior and Cognitive Behavioral Therapy for depression and/or anxiety
5. When starting pharmacotherapy, try to start only one at a time, if at all possible. Starting one medication at a time makes judging the effectiveness or adverse effects of that medication much more feasible.
6. When starting a new medication, try to assess symptom change by administering a pre and post symptom scale (e.g., Connors' Teachers Rating Scale or Teacher Report Form for ADHD symptoms, Overt Aggression Scale for Aggression, Young Mania Scale for bipolar disorder) or by quantifying the frequency and severity of target symptoms before and after the medication has been added.

7. If you inherit a patient who is already on multiple medications, try to understand the history of how your patient ended up on this regimen. Try to determine whether each medication has had positive, negative or no effects. If there is no evidence that a particular medication has been effective, try to discontinue it slowly and continue to monitor for increased symptoms. If an increase in symptoms is demonstrated, the medication can be re-started.
8. If a child or adolescent has done very well on a medication regimen for 12 months, consider the possibility of discontinuing one of his/her medications.

## **References**

- Jellinek, M. (1998). Overzealous prescribing of medications [Letter to the editor]. *J Am Acad Child Adolescent Psychiatry*, 37(9).
- Martin A, Van Hoof T, Stubbe D, Sherwin T, Scahill L. (2003). Multiple psychotropic pharmacotherapy among child and adolescent enrollees in Connecticut Medicaid managed care. *Psychiatric Services*, 54(1), 72-7.
- Martin A, Leslie D. (2003). Psychiatric inpatient, outpatient, and medication utilization and costs among privately insured youths, 1997-2000. *American Journal of Psychiatry*, 160(4): 757-64.
- National Association of State Mental Health Program Directors. (2001). Medical Directors' Technical Report on Psychiatric Polypharmacy.
- Olfson M, Marcus SC, Weissman MM, Jensen PS. (2000). National trends in the use of psychotropic medications by children. *J Am Academy Child Adolescent Psychiatry*, 41(5): 514-21
- Preda, A., et al. (1998). Premature psychopharmacology [Letter to the editor]. *J Am Academy Child Adolescent Psychiatry*, 37(4).
- Safer, D. (2003). Concomitant psychotropic medication for youths. *Am J Psychiatry*, 160, 438-449.
- Wakefield, T. (1999). The combined pharmacotherapy controversy [Letter to the editor]. *J Am Academy Child Adolescent Psychiatry*, 35(1).
- Woolston, J. (1999). Combined pharmacotherapy: Pitfalls of treatment. *J Am Academy Child Adolescent Psychiatry*, 38(11).

## **Other Useful Studies**

Arnold LE, Aman MG, Martin A, Collier-Crespin A, Vitiello B, Tierney E, Asarnow R, Bell-Bradshaw F, Freeman BJ, Gates-Ulanet P, Klin A, McCracken JT, McDougle CJ, McGough JJ, Posey DJ, Scahill L, Swiezy NB, Ritz L, Volkmar F. (2000). Assessment in multisite randomized clinical trials of patients with autistic disorder: the Autism RUPP Network. Research Units on Pediatric Psychopharmacology. *J Autism Dev Disorders*, Apr, (2): 99-111.

Jensen PS, Hinshaw SP, Swanson JM, Greenhill LL, Conners CK, Arnold LE, Abikoff HB, Elliott G, Hechtman L, Hoza B, March JS, Newcorn JH, Severe JB, Vitiello B, Wells K, Wigal T. (2001). Findings from the NIMH Multimodal Treatment Study of ADHD (MTA): implications and applications for primary care providers. *J Dev Behavioral Pediatrics*, 22(1), 60-73.

March JS, Swanson JM, Arnold LE, Hoza B, Conners CK, Hinshaw SP, Hechtman L, Kraemer HC, Greenhill LL, Abikoff HB, Elliott LG, Jensen PS, Newcorn JH, Vitiello B, Severe J, Wells KC, Pelham WE. (2000) Anxiety as a predictor and outcome variable in the multimodal treatment study of children with ADHD (MTA). *J Abnormal Child Psychology*, 28(6), 527-41.

(Fluvoxamine treatment, 2001). Fluvoxamine for the treatment of anxiety disorders in children and adolescents. The Research Unit on Pediatric Psychopharmacology Anxiety Study Group. *N England J Medicine*, 344(17), 1279-85.

(14-Month trial, 1999). A 14-month randomized clinical trial of treatment strategies for attention-deficit/hyperactivity disorder. The MTA Cooperative Group. Multimodal Treatment Study of Children with ADHD. *Arch Gen Psychiatry*, 56(12): 1073-86.

Pappadopulos E, Macintyre li JC, Crismon ML, Findling RL, Malone RP, Derivan A, Schooler N, Sikich L, Greenhill L, Schur SB, Felton CJ, Kranzler H, Rube DM, Sverd J, Finnerty M, Ketner S, Siennick SE, Jensen PS. (2003). Treatment recommendations for the use of antipsychotics for aggressive youth (TRAAY). Part II. *J Am Academy Child Adolescent Psychiatry*, 42(2), 145-61.

Schur SB, Sikich L, Findling RL, Malone RP, Crismon ML, Derivan A, Macintyre li JC, Pappadopulos E, Greenhill L, Schooler N, Van Orden K, Jensen PS. (2003). Treatment recommendations for the use of antipsychotics for aggressive youth (TRAAY). Part I: a review. *J Am Academy Child Adolescent Psychiatry*, 42(2), 132-44.

Tourette's Syndrome Study Group. (2002). Treatment of ADHD in children with tics: a randomized controlled trial. *Neurology*, 58(4), 527-36.

Vitiello B, Severe JB, Greenhill LL, Arnold LE, Abikoff HB, Bukstein OG, Elliott GR, Hechtman L, Jensen PS, Hinshaw SP, March JS, Newcorn JH, Swanson JM, Cantwell DP. (2001). Methylphenidate dosage for children with ADHD over time under controlled conditions: lessons from the MTA. *J Am Academy Child Adolescent Psychiatry*, 40(2), 188-96.

Walkup J, Labellarte M, Riddle MA, Pine DS, Greenhill L, Fairbanks J, Klein R, Davies M, Sweeney M, Abikoff H, Hack S, Klee B, Bergman RL, Lynn D, McCracken J, March J, Gammon P, Vitiello B, Ritz L, Roper M. (2002). Research Units on Pediatric Psychopharmacology Anxiety Study Group. Treatment of pediatric anxiety disorders: an open-label extension study. *J Child Adolescent Psychopharmacology*, 12(3), 175-88.

## **Reviews with Algorithms including Children's Medication Algorithm Project**

Birmaher B, Brent D, Heydl P. Childhood-onset depressive disorders. In: Fawcett J, Stein DJ, Jobson KO, eds. (1999). *Textbook of treatment algorithms in psychopharmacology*. New York: John Wiley & Sons, 135-163.

Davanzo PA. Mood stabilizers in the treatment of juvenile bipolar disorder: advances and controversies. (2000). *Child Adolescent Psychiatry Clin North Am*, 9(1),159-182.

Emslie GJ, Mayes TL, Hughes CW. Updates in the pharmacologic treatment of childhood depression. (2000). *Psychiatry Clin North Am Annual of Drug Therapy*, 7,235-256.

Hughes CW, Emslie GJ, Crismon ML, Wagner KD, Birmaher B, Geller B, Pliszka SR, Ryan ND, Strober M, Trivedi MH, Toprac MG, Sedillo A, Llana ME, Lopez M, Rush AJ. (1999). The Texas Children's Medication Algorithm Project: report of the Texas Consensus Conference Panel on Medication Treatment of Childhood Major Depressive Disorder. *J Am Academy Child Adolescent Psychiatry*, 38(11),1442-54. Latest update at <http://www.mhmr.state.tx.us/centraloffice/medicaldir>

Martin A, Kaufman J, Charney D. (2000). Pharmacotherapy of early-onset depression: update and new directions. *Child and Adolescent Psychiatry Clin North Am*, 9(1),135-157.

Pliszka SR, Greenhill LL, Crismon ML, Sedillo A, Carlson C, Conners CK, McCracken JT, Swanson JM, Hughes CW, Llana ME, Lopez M, Toprac MG. (2000). The Texas Children's Medication Algorithm Project: Report of the Texas Consensus Conference Panel on Medication Treatment of Childhood Attention-Deficit/Hyperactivity Disorder. Part II: Tactics. Attention-Deficit/Hyperactivity Disorder. *J Am Academy Child Adolescent Psychiatry*. 2000, 39(7),920-7.

Pliszka SR, Greenhill LL, Crismon ML, Sedillo A, Carlson C, Conners CK, McCracken JT, Swanson JM, Hughes CW, Llana ME, Lopez M, Toprac MG. (2000). The Texas Children's Medication Algorithm Project: Report of the Texas Consensus Conference Panel on Medication Treatment of Childhood Attention-Deficit/Hyperactivity Disorder. Part I. Attention-Deficit/Hyperactivity Disorder. *J Am Academy Child Adolescent Psychiatry*,39(7),908-19.Update: <http://www.mhmr.state.tx.us/centraloffice/medicaldir>

Hughes CW, Emslie GJ, Crismon ML, Wagner KD, Birmaher B, Geller B, Pliszka SR, Ryan ND, Strober M, Trivedi MH, Toprac MG, Sedillo A, Llana ME, Lopez M, Rush AJ. (1999). The Texas Children's Medication Algorithm Project: report of the Texas Consensus Conference Panel on Medication Treatment of Childhood Major Depressive Disorder. *J Am Academy Child Adolescent Psychiatry*, 38(11),1442-54. Latest update at <http://www.mhmr.state.tx.us/centraloffice/medicaldir>

## **Practice Guidelines**

American Academy of Pediatrics. Subcommittee on Attention-Deficit/Hyperactivity Disorder and Committee on Quality Improvement. (2001). Clinical practice guideline: treatment of the school-aged child with attention-deficit/hyperactivity disorder. *Pediatrics*, 108(4),1033-44. \* Atomoxetine not included

Dulcan M. (1997). Practice parameters for the assessment and treatment of children, adolescents, and adults with attention-deficit/hyperactivity disorder. American Academy of Child and Adolescent Psychiatry. *J Am Academy Child Adolescent Psychiatry*, 36(Suppl 10), 85S-121S. \*Atomoxetine not included

King RA. (1997). Practice parameters for the psychiatric assessment of children and adolescents. American Academy of Child and Adolescent Psychiatry. *J Am Academy Child Adolescent Psychiatry*, 36(Suppl 10),4S-20S.

(Practice parameters posttraumatic, 1998). Practice parameters for the assessment and treatment of children and adolescents with posttraumatic stress disorder. *J Am Academy Child Adolescent Psychiatry*, 37(Suppl 10), 4S-26S.

(Practice parameters OCD,1998). Practice parameters for the assessment and treatment of children and adolescents with obsessive-compulsive disorder. AACAP. *J Am Academy Child Adolescent Psychiatry*, 37(Suppl 10),27S-45S.

(Guideline ADHD, 2000). Clinical practice guideline: diagnosis and evaluation of the child with attention-deficit/hyperactivity disorder. *Am Academy Pediatrics*, 105(5),1158-70. \*Atomoxetine not included

(Practice parameters depressive disorders, 1998). Practice parameters for the assessment and treatment of children and adolescents with depressive disorders. AACAP. *J Am Academy Child Adolescent Psychiatry*, 37(Suppl 10),63S-83S. \*Written before Dear Professional letter about paroxetine sent by FDA

(Practice parameters bipolar disorder, 1997). AACAP official action. Practice parameters for the assessment and treatment of children and adolescents with bipolar disorder. *J Am Academy Child Adolescent Psychiatry*, 36(1),138-57.

Szymanski L, King BH. (1999). Practice parameters for the assessment and treatment of children, adolescents, and adults with mental retardation and comorbid mental disorders. American Academy of Child and Adolescent Psychiatry Working Group on Quality Issues. *J Am Academy Child Adolescent Psychiatry*, 38(12 Suppl),5S-31S.

Thomas JM, Benham AL, Gean M, Luby J, Minde K, Turner S, Wright HH. (1997). Practice parameters for the psychiatric assessment of infants and toddlers (0-36 months). American Academy of Child and Adolescent Psychiatry. *J Am Academy Child Adolescent Psychiatry*, 36(Suppl 10), 21S-36S.

Volkmar F, Cook EH Jr, Pomeroy J, Realmuto G, Tanguay P. (1999). Practice parameters for the assessment and treatment of children, adolescents, and adults with autism and other pervasive developmental disorders. American Academy of Child and Adolescent Psychiatry Working Group on Quality Issues. *J Am Academy Child Adolescent Psychiatry*, 38(Suppl 12), 32S-54S. Erratum in: *J Am Academy Child Adolescent Psychiatry* 2000 Jul;39(7):938.